

Extensions of Finite Abelian Groups

Guhan Venkat

February 12, 2010

Abstract

We study group extensions of Finite Abelian Groups using matrices. We also prove a Theorem for equivalence of extensions using matrices.

Introduction

Presentation of Abelian Groups

Every Abelian Group $(A, +)$ can be considered as a $\mathbb{Z}$ -module under the ring action:

$$\begin{aligned}\mathbb{Z} \times A &\longrightarrow A \\ (n, x) &\longrightarrow nx = x + x + \dots + x (n \text{ times}) \\ (-n, x) &\longrightarrow -(nx) = -(x + x + x + \dots + x) (n \text{ times})\end{aligned}$$

We call $\vec{x}^t = (x_1, x_2, \dots, x_n), x_i \in A$ a generating set of A if every element in A can be written as $a = \alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n$ for some $\alpha_i \in \mathbb{Z}$ under the ring action defined above.

A relation in A is an equation of the form

$$\alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n = 0$$

If we have a set of 'm' relations

$$\alpha_{11}x_1 + \alpha_{12}x_2 + \dots + \alpha_{1n}x_n = 0$$

$$\alpha_{21}x_1 + \alpha_{22}x_2 + \dots + \alpha_{2n}x_n = 0$$

$\vdots$

$$\alpha_{n1}x_1 + \alpha_{n2}x_2 + \dots + \alpha_{nn}x_n = 0$$

We can write that down as a matrix equation $\alpha \vec{x} = 0$ or

$$\begin{pmatrix} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & \alpha_{nm} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = 0$$

A complete set of relations of a group A is one from which every other relation of the group can be derived by a linear combination. A 'Presentation of an Abelian Group' is a generating set along with a complete set of relations.

Finite Abelian Groups and their Sylow Subgroups

Let A be a Finite Abelian Groups of order n where $n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_m^{\alpha_m}$ then $A \cong A_{p_1} \times A_{p_2} \times \dots \times A_{p_m}$ where each A_{p_i} is a Sylow subgroup associated with that prime. Therefore, we study abelian p-groups.

The Extension Problem for abelian p-groups

Given two p-groups G_λ and G_μ of type λ and type μ respectively, we describe all groups E containing G_λ such that $E/G_\lambda \cong G_\mu$.

i.e. we have a Short Exact Sequence $0 \longrightarrow G_\lambda \longrightarrow E \longrightarrow G_\mu \longrightarrow 0$

By the structure theorem for finite abelian groups, G_λ has a presentation with generating set $y = \langle y_1, y_2, \dots, y_l \rangle$ of l elements and relations $p^{\lambda_i} y_i = 0 \forall i$ from 1 to l and for some integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l > 0$

Similarly G_μ is generated by m elements $\langle x_1, x_2, \dots, x_m \rangle$ with the relations $p^{\mu_j} x_j = 0 \forall j$ from 1 to m

Now choose $x_1^\sim, x_2^\sim, x_3^\sim, \dots, x_m^\sim \in E$ such that $\phi(x_j^\sim) = x_j \forall j$ from 1 to m

But from the above relations, we have $\phi(p^{\mu_j} x_j^\sim) = p^{\mu_j} \phi(x_j^\sim) = p^{\mu_j} x_j = 0 \Rightarrow p^{\mu_j} x_j^\sim \in \ker(\phi) = G_\lambda$ (Definition of Short Exact Sequence)

Since every element of G_λ can be written as some $a \cdot \vec{y}$ we have $p^{\mu_j} x_j^\sim = a_{j1} y_1 + a_{j2} y_2 + \dots + a_{jl} y_l \forall j$ from 1 to m

Combining the two sets of relations, we have in hand altogether $l + m$ relations. Putting these relations together as a matrix, we would get a pre-

sentation for E

$$\begin{pmatrix} p^{\lambda_1} & & & & & & & \\ & p^{\lambda_2} & & & & & & \\ & & \ddots & & & & & \\ & & & p^{\lambda_l} & & & & \\ a_{11} & a_{12} & \dots & a_{1l} & p^{\mu_1} & & & \\ a_{21} & a_{22} & \dots & a_{2l} & & p^{\mu_2} & & \\ \vdots & & & & & & \ddots & \\ a_{21} & a_{22} & \dots & a_{ml} & \dots & \dots & \dots & p^{\mu_m} \end{pmatrix}_{m+l \times m+l} \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_l \\ x_1^\sim \\ x_2^\sim \\ \vdots \\ x_m^\sim \end{pmatrix} = 0$$

Or as blocks they would be

$$\begin{pmatrix} p^\lambda & \\ A & p^\mu \end{pmatrix}_{m+l \times m+l} \begin{pmatrix} y \\ x^\sim \end{pmatrix} = 0$$

One is now interested in classifying extensions upto isomorphisms

$$\begin{array}{ccccccc} 0 & \rightarrow & G_\lambda & \xrightarrow{\psi_1} & E_1 & \xrightarrow{\phi_1} & G_\mu \rightarrow 0 \\ & & \downarrow f & & \downarrow h & & \downarrow g \\ 0 & \rightarrow & G'_\lambda & \xrightarrow{\psi_2} & E_2 & \xrightarrow{\phi_2} & G'_\mu \rightarrow 0 \end{array}$$

If there exist isomorphisms f and g and a homomorphism h such that the above diagram commutes, then the extension in the first row is said to be equivalent to the extension in the second row. It then follows that h is also an isomorphism.

$$G'_\lambda = \langle y'_i \rangle$$

where i is from 1 to l

$$G'_\mu = \langle x'_j \rangle$$

where j is from 1 to m (We have assumed that isomorphic groups have equal number of generators and relations. In rigor, one should look at isomorphic groups through Tietze Transformations.)

We can now describe isomorphisms f, g and h as matrices acting on basis (vectors).

$$f(\langle y_1, y_2, \dots, y_l \rangle) = \langle y'_1, y'_2, \dots, y'_l \rangle [F]$$

$$g(\langle x_1, x_2, \dots, x_m \rangle) = \langle x'_1, x'_2, \dots, x'_m \rangle [G]$$

$$h(\langle y_1, \dots, y_l, x_1^\sim, \dots, x_m^\sim \rangle) = \langle y'_1, \dots, y'_l, x_1^{\sim'}, \dots, x_m^{\sim'} \rangle [H]$$

Here $[F]$, $[G]$ and $[H]$ are the matrices of automorphisms. We write $[H]$ as

$$\begin{bmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{bmatrix} \text{ where } h_{11} \text{ is } l \times l \text{ and } h_{22} \text{ is } m \times m.$$

Also the matrices are invertible and preserve the structure of the groups, i.e

$$\begin{aligned}
F &\in GL_l(\mathbb{Z}) \cap p^{-\lambda} GL_l(\mathbb{Z}) p^\lambda \\
G &\in GL_m(\mathbb{Z}) \cap p^{-\mu} GL_m(\mathbb{Z}) p^\mu \\
H &\in GL_{l+m}(\mathbb{Z}) \cap p^{-(\lambda+\mu)} GL_{l+m}(\mathbb{Z}) p^{(\lambda+\mu)}
\end{aligned}$$

Using the fact that the above diagram commutes, we have

$$\psi_2(f(k)) = h(\psi_1(k)) \forall k \in G_\lambda$$

$$\begin{aligned}
\begin{bmatrix} I \\ 0 \end{bmatrix} [F] &= \begin{bmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{bmatrix} \begin{bmatrix} I \\ 0 \end{bmatrix} \\
\Rightarrow \begin{bmatrix} F \\ 0 \end{bmatrix} &= \begin{bmatrix} h_{11} \\ h_{21} \end{bmatrix} \Rightarrow h_{11} = F \text{ and } h_{21} = 0
\end{aligned}$$

$$\phi_2(h(l)) = g(\phi_1(l)) \forall l \in E_1$$

$$\begin{aligned}
\begin{bmatrix} F & h_{12} \\ 0 & h_{22} \end{bmatrix} \begin{bmatrix} 0 \\ I \end{bmatrix} &= \begin{bmatrix} 0 \\ I \end{bmatrix} [G] \\
\Rightarrow \begin{bmatrix} h_{12} \\ h_{22} \end{bmatrix} &= \begin{bmatrix} 0 \\ G \end{bmatrix} \\
\Rightarrow [H] &= \begin{bmatrix} F & 0 \\ 0 & G \end{bmatrix}
\end{aligned}$$

Now, looking at the presentations for E_1 and E_2

$$E_1 : \begin{bmatrix} y & x^\sim \end{bmatrix} \begin{bmatrix} A_1 & p^\lambda \\ p^\mu & 0 \end{bmatrix} = 0$$

$$E_1 : \begin{bmatrix} y' & x'^\sim \end{bmatrix} \begin{bmatrix} A_2 & p^\lambda \\ p^\mu & 0 \end{bmatrix} = 0$$

Note that the matrices have been transposed for convinience.

Using the isomorphism condition, i.e. $h(0) = 0$, we have

$$\begin{aligned}
\begin{bmatrix} y & x^\sim \end{bmatrix} \begin{bmatrix} F & 0 \\ 0 & G \end{bmatrix} \begin{bmatrix} A_1 & p^\lambda \\ p^\mu & 0 \end{bmatrix} &= 0_{E_2} \\
\begin{bmatrix} y & x^\sim \end{bmatrix} \begin{bmatrix} FA_1 & Fp^\lambda \\ Gp^\mu & 0 \end{bmatrix} &= 0_{E_2}
\end{aligned}$$

But given that we have a complete set of relations in E_2 , these matrices must be related by a column operation (Note, we had transposed the matrices and hence instead of row operations, we now must consider column operations).

$$\begin{bmatrix} A_2 & p^\lambda \\ p^\mu & 0 \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} = \begin{bmatrix} FA_1 & Fp^\lambda \\ Gp^\mu & 0 \end{bmatrix}$$

$$\begin{aligned}
&\Rightarrow p^\mu \alpha = Gp^\mu \text{ or } \alpha = p^{-\mu} Gp^\mu \\
&\Rightarrow A_2 p^{-\mu} Gp^\mu + p^\lambda \gamma = F A_1 \\
&\Rightarrow F^{-1} A_2 p^{-\mu} Gp^\mu + F^{-1} p^\lambda \gamma = A_1
\end{aligned}$$

Now, given the conditions on F and G , we re-write the above equation as

$$F^{-1} A_2 G' + F^{-1} p^\lambda \gamma = A_1$$

But, A_1 is taken modulo $p^\lambda \Rightarrow F' A_2 G' = A_1$ where $G' = p^{-\mu} Gp^\mu$ and $F' = F^{-1}$. Thus we have the following Theorem.

Theorem The extensions associated to two finite abelian groups E_1 and E_2 are equivalent if and only if there exist matrices $F \in GL_l(\mathbb{Z}) \cap p^\lambda GL_l(\mathbb{Z}) p^{-\lambda}$ and $G \in GL_m(\mathbb{Z}) \cap p^{-\mu} GL_m(\mathbb{Z}) p^\mu$ such that $F A_2 G = A_1$ where the (i, j) entry of each matrix is taken modulo $p^{\min(\lambda_i, \mu_j)}$

Acknowledgements

This project was carried out under the guidance of Dr. Amritanshu Prasad, IMSc as a part of IAS - SRFP 2009.

References

- [1] Garrett Birkhoff. Subgroups of Abelian Groups *Proc. London Math. Soc.* 38:385-401, 1935.
- [2] David S. Dummit and Richard M. Foote *Abstract Algebra Third Edition* John Wiley and Sons inc. 2004.